arXiv:1408.2364v3 [cs.CC] 17 Nov 2014

Notes on space complexity of integration of computable real functions in Ko–Friedman model

Sergey V. Yakhontov

Abstract

In the present paper it is shown that real function $g(x) = \int_0^x f(t)dt$ is a linear-space computable real function on interval $[0,1]$ if f is a linear-space computable $C^2[0,1]$ real function on interval $[0,1]$, and this result does not depend on any open question in the computational complexity theory. The time complexity of computable real functions and integration of computable real functions is considered in the context of Ko–Friedman model which is based on the notion of Cauchy functions computable by Turing machines.

In addition, a real computable function f is given such that $\int_0^1 f \in \mathbf{FDSPACE}(n^2)_{C[a,b]}$ but $\int_0^1 f \notin \mathbf{FP}_{C[a,b]}$ if $\mathbf{FP} \neq \#\mathbf{P}$.

Keywords: Computable real functions, Cauchy function representation, polynomial-time computable real functions, linear-space computable real functions, $C^2[0,1]$ real functions, integration of computable real functions.

Contents

1 Introduction **1**
1.1 CF computable real numbers and functions 2
1.2 Integration of **FP** computable real functions 2

2 Upper bound of the time complexity of integration **3**

3 Function from $\mathbf{FDSPACE}(n^2)_{C[a,b]}$ that not in $\mathbf{FP}_{C[a,b]}$ if $\mathbf{FP} \neq \#\mathbf{P}$ **4**

4 Conclusion **4**

1 Introduction

In the present paper, we consider computable real numbers and functions that are represented by Cauchy functions computable by Turing machines [1].

Main results regarding computable real numbers and functions can be found in [1–4]; main results regarding computational complexity of computations on Turing machines can be found in [5].

As is usual, the set of real functions whose 2-nd derivative exists and is continuous on interval $[0,1]$ is denoted by $C^2[0,1]$, and the set of $C^k[0,1]$ real functions for all $k \geq 1$ is denoted by $C^\infty[0,1]$.

Sergey V. Yakhontov: Ph.D. in Theoretical Computer Science, Dept. of Computer Science, Faculty of Mathematics and Mechanics, Saint Petersburg State University, Saint Petersburg, Russian Federation, 198504;
e-mail: SergeyV.Yakhontov@gmail.com, S.Yakhontov@spbu.ru; phone: +7-911-966-84-30;
personal Web page: https://sites.google.com/site/sergeyvyakhontov/;17-Nov-14

It is known [1] that real function $g(x) = \int_0^x f(t)dt$ is polynomial-time computable real function on interval $[0,1]$ iff $\mathbf{FP} = \mathbf{\#P}$ wherein f is a polynomial-time computable real function on interval $[0,1]$. It means integration of polynomial-time computable real functions is as hard as string functions from complexity class $\mathbf{\#P}$.

So this result from [1] is relativized to question whether $\mathbf{FP} = \mathbf{\#P}$ or not which is one of the open questions in the computational complexity theory.

In the present paper it is shown that real function $g(x) = \int_0^x f(t)dt$ is a linear-space computable real function on interval $[0,1]$ if f is a linear-space computable $C^2[0,1]$ real function on interval $[0,1]$, and this result does not depend on any open question in the computational complexity theory.

1.1 CF computable real numbers and functions

Cauchy functions in the model defined in [1] are functions binary converging to real numbers. A function $\phi : \mathbb{N} \to \mathbf{D}$ (here $\mathbf{D}$ is the set of dyadic rational numbers) is said to binary converge to real number x if

$$|\phi(n) - x| \leq 2^{-n}$$

for all $n \in \mathbb{N}$; CF_x denotes the set of all functions binary converging to x.

Real number x is said to be a CF computable real number if CF_x contains a computable function ϕ.

Real function f on interval $[a,b]$ is said to be a CF computable function on interval $[a,b]$ if there exists a function-oracle Turing machine M such that for all $x \in [a,b]$ and for all $\phi \in CF_x$ function ψ computed by M with oracle ϕ is in $CF_{f(x)}$.

The input of functions ϕ and ψ is 0^n (0 repeated n times) when a number or a function is evaluated to precision 2^{-n}.

Definition 1. [1] *Function $f : [a,b] \to \mathbb{R}$ is said to be computable in time $t(n)$ real function on interval $[a,b]$ if for all computable real numbers $x \in [a,b]$ function $\psi \in CF_{f(x)}$ (ψ is from the definition of CF computable real function) is computable in time $t(n)$.*

Definition 2. [1] *Function $f : [a,b] \to \mathbb{R}$ is said to be computable in space $s(n)$ real function on interval $[a,b]$ if for all computable real numbers $x \in [a,b]$ function $\psi \in CF_{f(x)}$ (ψ is from the definition of CF computable real function)* is computable in space $s(n)$.

FP denotes the class of string functions computable in polynomial time on Turing machines, **FLINSPACE** denotes the class of string functions computable in linear space on Turing machines, and **FEXPTIME** denotes the class of string functions computable in exponential time on Turing machines.

According to these notations, polynomial-time computable real functions are said to be **FP** computable real functions, linear-space computable real functions are said to be **FLINSPACE** computable real functions, and exponential-time computable real functions are said to be **FEXPTIME** computable real functions.

The set of **FP** computable real functions on interval $[a,b]$ is denoted by $\mathbf{FP}_{C[a,b]}$, the set of **FLINSPACE** computable real functions on interval $[a,b]$ is denoted by $\mathbf{FLINSPACE}_{C[a,b]}$, and the set of **FEXPTIME** computable real functions on interval $[a,b]$ is denoted by $\mathbf{FEXPTIME}_{C[a,b]}$.

The set of the $C^2[0,1]$ real functions from class $\mathbf{FP}_{C[a,b]}$ is denoted by $\mathbf{FP}_{C^2[a,b]}$, and the set of the $C^\infty[0,1]$ real functions from class $\mathbf{FP}_{C[a,b]}$ is denoted by $\mathbf{FP}_{C^\infty[a,b]}$.

The same definitions are for other complexity classes.

1.2 Integration of FP computable real functions

The main results from [1] regarding integration of **FP** computable real functions are the following.

Theorem 1. [1, 5.33] *The following are equivalent:*

a) Let f be in $\mathbf{FP}_{C[0,1]}$. Then, the function $g(x) = \int_0^x f(t)dt$ is polynomial-time computable.

b) Let f be in $\mathbf{FP}_{C^\infty[0,1]}$. Then, the function $g(x) = \int_0^x f(t)dt$ is polynomial-time computable.

c) $\mathbf{FP} = \#\mathbf{P}$.

It means if $\mathbf{FP} \neq \#\mathbf{P}$ then the integral of a polynomial-time computable real function f is not necessarily polynomial-time computable even if f is known to be infinitely differentiable. But if f is polynomial-time computable and is analytic on $[0, 1]$ then the integral of f must be computable in polynomial time.

Some additional results regarding the time complexity of integration of computable real functions can be found in [6]. For example, the computation of the volume of a one-dimensional convex set K is $\#\mathbf{P}$-complete if K is represented by a polynomial-time computable function defining its boundary.

2 Upper bound of the time complexity of integration

Let f be a linear-space computable $C^2[0, 1]$ real function on interval $[0, 1]$. Let's consider the composite trapezoidal rule [7] for function f on interval $[a, x]$ for $a < x \leq b$:

$$\begin{aligned} g(x) &= \int_a^x f(t)dt = \\ &= \frac{h}{2}\left(f(t_0) + \left(\sum_{i=1}^{k-1} 2 \cdot f(t_i)\right) + f(t_k)\right) - \\ &\quad - \frac{(b-a)^3}{12 \cdot k^2} f''(\xi) \end{aligned}$$

wherein k is a natural number, $h = \frac{x-a}{k}$, $t_0 = a$, $t_i = a + i \cdot h$, $t_k = b$, and $a < \xi < x$. This equation on interval $[0, x]$ is as follows:

$$\begin{aligned} g(x) &= \int_0^x f(t)dt = \\ &= \frac{h}{2}\left(f(t_0) + \left(\sum_{i=1}^{k-1} 2 \cdot f(t_i)\right) + f(t_k)\right) - \\ &\quad - \frac{1}{12 \cdot k^2} f''(\xi) \end{aligned} \tag{1}$$

wherein $h = \frac{x}{k}$, $t_0 = 0$, $t_i = i \cdot h$, $t_k = x$, and $0 < \xi < x$.

To compute approximations $g^*(x)$ of function g on interval $[0, 1]$, let's compute approximations $f^*(x_i)$ of function f at points x_i, $i \in [0..k]$, to precision 2^{-m} wherein m is a natural number. In that case, the following holds:

$$\begin{aligned} |\Delta(g; x)| &= |g^*(x) - g(x)| = \\ &= \left|\frac{h}{2}\left(f^*(t_0) + \left(\sum_{i=1}^{k-1} 2 \cdot f^*(t_i)\right) + f^*(t_k)\right) - \right. \\ &\quad \left. - \frac{h}{2}\left(f(t_0) + \left(\sum_{i=1}^{k-1} 2 \cdot f(t_i)\right) + f(t_k)\right)\right| + \\ &\quad + \frac{1}{12 \cdot k^2} f''(\xi) \leq \\ &\leq \frac{h}{2}\left(|f^*(t_0) - f(t_0)| + \left(\sum_{i=1}^{k-1} 2 \cdot |f^*(t_i) - f(t_i)|\right) + \right. \\ &\quad \left. + |f^*(t_n) - f(t_k)|\right) + \frac{1}{12 \cdot k^2} f''(\xi). \end{aligned}$$

So, we have:

$$|\Delta(g;x)| = |g^*(x) - g(x)| \leq$$
$$\leq \frac{h}{2}\left(\frac{1}{2^m} + \frac{2(k-1)}{2^m} + \frac{1}{2^m}\right) + \frac{1}{12\cdot k^2}C_1 \leq$$
$$\leq \frac{1}{k}\frac{k}{2^m} + \frac{1}{12\cdot k^2}C_1 = \frac{1}{2^m} + \frac{1}{k^2}C_2;$$

here the fact that f'' is bounded above if $f \in C^2[0,1]$ is taken into account. If we take $m = 2n$ and $k = C_2 2^n$ then

$$|\Delta(g;x)| = \frac{1}{2^{2n}} + \frac{1}{(C_2)^2 2^{2n}}C_2 < 2^{-n}.$$

It means it is sufficient to compute in a loop for $i \in [0,k]$ approximations $g^*(x)$ of function g on interval $[0,1]$ using formula (1) wherein $k = C_2 2^n$ and $f^*(x_i)$ are the approximations of function f at points x_i, $i \in [0..k]$, to precision 2^{-m} wherein $m = 2n$.

So, the following theorems holds.

Theorem 2. *If f is a linear-space computable $C^2[0,1]$ real function on interval $[0,1]$ then real function $g(x) = \int_0^x f(t)dt$ is a linear-space computable real function on interval $[0,1]$.*

The same holds for class $\mathbf{FDSPACE}(n^2)$.

Theorem 3. *If f is a $\mathbf{FDSPACE}(n^2)$ computable $C^2[0,1]$ real function on interval $[0,1]$ then real function $g(x) = \int_0^x f(t)dt$ is a $\mathbf{FDSPACE}(n^2)$ computable real function on interval $[0,1]$.*

Theorem 4. *If f is a polynomial-time and linear-space computable $C^2[0,1]$ real function on interval $[0,1]$ then real function $g(x) = \int_0^x f(t)dt$ is an exponential-time and linear-space computable real function on interval $[0,1]$.*

3 Function from $\mathbf{FDSPACE}(n^2)_{C[a,b]}$ that not in $\mathbf{FP}_{C[a,b]}$ if $\mathbf{FP} \neq \#\mathbf{P}$

Let's consider fucntion $f(x) = \sum_{n=1}^{\infty} f_n(x)$ from proof $(d) \Rightarrow (e)$ of theorem 5.32 [1, p.186]. Some of the properties of function f are as follows:

1) $f \in C^\infty[0,1]$;
2) if $\int_0^1 f \in \mathbf{FP}_{C[0,1]}$ then $\mathbf{FP} = \#\mathbf{P}$;
3) $f \in \mathbf{FDSPACE}(n^2)_{C[a,b]}$ if one takes set B ([1, theorem 5.32, p.184]) such that $B \in \mathbf{DSPACE}(n^2)$;
4) $\int_0^1 f \in \mathbf{FDSPACE}(n^2)_{C[a,b]}$ according to theorem 3.

Therefore, the following theorem holds

Theorem 5. $\mathbf{FP}_{C[0,1]} \neq \mathbf{FDSPACE}(n^2)_{C[a,b]}$ *if* $\mathbf{FP} \neq \#\mathbf{P}$.

4 Conclusion

In the present paper, it is shown that real function $g(x) = \int_0^x f(t)dt$ is a linear-space computable real function on interval $[0,1]$ whenever f is a linear-space computable $C^2[0,1]$ real function on interval $[0,1]$. This result differs from the result from [1] regarding the time complexity of integration of polynomial-time computable real functions in the sense that integration of a polynomial-time computable real function **may** be not polynomial-time computable (if f is not analytic and $\mathbf{FP} \neq \#\mathbf{P}$), but integration of linear-space computable C^2 real functions is **always** linear-space computable.

Regarding further investigations, it is interesting to derive results regarding the space complexity of other operators, for example, the space complexity of differentiation of computable functions.

References

[1] Ko K. *Complexity Theory of Real Functions.* Boston: Birkhauser, 1991. 309 p.

[2] Aberth O. *Computable calculus.* Academic Press, 2001. 192 p.

[3] Kushner B. A. *Lectures on Constructive Mathematical Analysis.* American Mathematical Society, 1984. 346 p.

[4] Weihrauch K. *Computable analysis.* New York: Springer, 2000. 285 p.

[5] Du D., Ko K. *Theory of Computational Complexity.* New York: John Wiley & Sons, 2000. 491 p.

[6] Du D., Ko K. *Computational complexity of integration and differentiation on convex functions* // System Sci. and Math. Sci 2, pp. 70–79.

[7] Berezin I. S., Zhidkov N. P. *Numeric analysis.* M.: Fiz-Mat Lit, 1962, Vol. 1. 464 p.